

2. LACUSTRIS.

Animal nigrescens. Tentacula breviora.

Testa oblonga, subdepressa, vertice excentrico; fragilis, membranacea, concentricè substriata, nigrescens.

Apertura oblonga.

Long. 0.1.—Diam. 0.35.

Ancylus lacustris. Müll. Verm. 2. p. 199. Drap. Hist. des Moll. p. 47. t. 2. f. 25—27.

Patella oblonga. Lightf. in Phil. Trans. 76. p. 168. t. 3. f. 1—5.

——— *lacustris. Gmel. Syst. Nat. 1. p. 3710. Mont. Test. Brit. p. 484.*

On water-plants in ditches, lakes, &c.; though rather a local species. It is not uncommon in Blackpill marsh near Swansea, mixed with the *Planorbis nitidus*.

 ADDITIONS AND CORRECTIONS.

Page 324, for *Cianella*, read *Cionella*.

HELIX FUSCA, p. 330.

Animal lutescenti-griseum, dorso flavescenti. Tentacula prælonga, flexilia.

Since writing the above list, I have discovered this local species in great abundance in moist parts of Penllergare wood, near Swansea, on the leaves and stems of the *Hera-cleum Sphondylium* and other plants.

HELIX NITIDA, p. 339.

The variety β . is found not uncommonly under leaves, decaying *Boleti*, &c. in the woods at Penllergare.

LIMNEUS TINCTUS, p. 378.

Is the young of a small variety of *communis*; but so very dissimilar in its form from that shell, that I could not, till after a careful examination, and comparison of a series of specimens of both species, be brought to unite them.